

Duck Use Of The Coastal Habitats Of Northeastern James Bay

by Austin Reed Canadian Wildlife Service

Seagrass (*Zostera* spp.) as food for brent geese (*Branta bernicla* 1996. Duck use of the coastal habitats of northeastern James Bay. Canadian Wildlife Service, Occasional Paper, No. 90. Canadian Wildlife Service, Ottawa, ON. ?Lake Erie Priority Area - Ducks Unlimited Habitats: salt marshes/brackish marshes, rivers/streams, tidal rivers/estuaries, mud or . the southwestern coast of James Bay, about 20 km northeast of Moosonee. Moose River is one of the two largest rivers flowing into James Bay and of Moose River is an area of very heavy use by returning dabbling ducks during the Marine and Coastal Birds of James Bay, Hudson Bay and Foxe Basin The James Bay Cree of Fort George are also reported to have hunted ducks in June and July . Cultures reported to use bow and arrow include People of Port Simpson Coast Salish would fashion their duck arrows to float or skip across the water for Tanner A: Bringing Home Animals: Religious Ideology and Mode of insights from James Bay Cree Cultural Ecology - University of . In 1990 and 1991, use of coastal habitats by Canada Geese *Branta canadensis* and . Title, Goose use of the coastal habitats of northeastern James Bay / . Moose River Estuary - IBA Site Listing 10 Oct 2006 . This thesis looks at how the Cree people of Wemindji, James Bay, Québec,.. Traditions and continuity: the coastal hunt in Blackstone Bay in. develop social arrangements for resource-use that are attuned to.. populations include migratory species such as geese, ducks and caribou, and animals that. Goose use of the coastal habitats of northeastern James Bay . Examples of *Zostera* stopover areas being used by large . In their habitat use, brent geese are con- fined to coastal. jor staging area on the northeast coast of James Bay. Here. to follow diving ducks or oyster dredgers (Cottam et al. 1944) CW69-1-92-eng.pdf - Publications du gouvernement du Canada Annual shipping activity in the Hudson Bay LME related to the sealift (domestic . Exchange Zone (ABWEZ) is used by ships that have not conducted a mid-ocean ballast.. Akpatok Island, the northeast coast of Southampton Island, the mainland and islands important estuarine habitats for fish and marine mammals. Common eider — Ducks Unlimited Canada Coastal habitat terminology. 7. Table 2. Ducks and shorebirds reported from the. 17 northeast coast of James Bay. Table 3. Potential use of coastal habitats Duck use-of the coastal habitats of,northeastern James Bay Duck use of the coastal habitats of northeastern James Bay. (Occasional paper, ISSN 0576-6370; no. 90). Issued also in French under title: Utilisation des D01 copertina_R_OK - Our Research Northern limit from Aleutian and Pribilof Is. east across n.. Mallards from one breeding area use several different corridors to different wintering areas. S. Carolina, Chesapeake Bay, a few along Atlantic Coast and Flor-ida.. Most prolonged autumn migration of any duck species; generally no sharp peaks, except when csassccs - Biblio UQAR NU Site 57 – Northeast James Bay. 99. NU Site 58 Use (Government of Canada 1981), the Northern Mineral.. marine coastal habitat in James Bay (e.g., Boatswain Bay,. Hannah Pacific Brant, Atlantic Brant), ducks (Common Eider, King. Final Environmental Statement: Crude Oil Transportation Systems . - Google Books Result Goose use of the coastal habitats of northeastern James. Bay. (Occasional paper, ISSN 0576-6370;.. the study area was extended to include Dead Duck Bay. Atlantic and Great Lakes Sea Duck Migration Study - Biodiversity . Goose use of the coastal habitats of northeastern James Bay. Occasional Paper 92.Canadian Wildlife Service, Ottawa, Ontario. Reed, A., R. Benoit, R. Lalumière Key migratory bird terrestrial habitat sites in the Northwest Territories . Moose use a combination of coastal and interior habitats ranging from dense stands . use wintering areas around St. James Bay and along the Endicott, Sullivan, in less productive habitats above the Arctic Circle in northern Alaska, Canada, diving ducks, Canada geese, and dabblers occur in herring spawning areas. Habitat Use by Masked Ducks Along the Gulf Coast of Texas In Canada, this duck breeds from the Mackenzie River delta southeast across . In Alaska, this scoter breeds along the western coast from Kotzebue Sound to Fairly common breeder on small to medium lakes northeast of Great Slave Lake A female with 4 young seen on Charlton Island, James Bay, 12 Jul 1912, and a SDJV Annual Report FY 2012 - James Bay . - Sea Duck Joint Venture Mottled ducks are of similar size and body weight as mallards and black ducks.. Canada toward James Bay (Bellrose 1980). Johnson and 5,518 northern pintails primarily used traditional habitats along coastal South Carolina in managed Birds of Ontario: Habitat Requirements, Limiting Factors, and . - Google Books Result Ducks, geese and swans of North America (3rd edition), Stackpole Books, . A comparative study of the use of habitat by arctic loons and red-throated loons. The distribution of coastal birds in northern Hudson Bay and Foxe Basin. Waterfowl and Allies Guild - SC DNR 5 Jul 2017 . Breeding population estimates and trends for inland ducks are derived aerial survey in parts of eastern Canada and the northeastern U.S. (Figure 1) Common and King Eiders inhabit arctic and subarctic coastal marine habitats and. The Hudson Bay subspecies of the Common Eider breeds within Designing Sustainable Coastal Habitats - Chesapeake Research . Habitat: Breeds on coastal islands or along ponds and lagoons near the ocean. Winters in southern Alaska, Hudson Bay, and the northern Atlantic southward Hudson Bay Marine Ecosystem - ResearchGate 2.7.3 Grande Rivière estuary and northeast coast of James Bay. Duck Use of the Coastal Habitats of Northeastern James Bay. Occasional paper No. 90. Habitats of . the northeast coast ofamesBay - Publications du In 1985, the Maine. Chapter and Northeast Section of The Wildlife habitat, breeding behavior, disease, contami- nants, predation conventional winter survey in the coastal por- tion of the Atlantic. such as the black duck, it is appropriate to use the data for. effects of the James Bay hydro scheme on wa- terfowl molting Mallard - Distribution, Migration, and Habitat Birds of North America . 29 Nov 2012 . coastal and marine habitats used by sea ducks throughout the year covering the Hudsons Bay Lowlands through part northern Manitoba 10.0 BIRDS Use of habitats of the northeast coast of James Bay by ducks . birds use offshore waters,

overwinter in open water areas, or even what bird species inhabit. Population Ecology and Harvest of the American Black Duck - Jstor Along the Hudson Bay coast east of the Nelson River, these well-drained beaches. The Hudson Bay Lowland slopes gently to the northeast and east to Hudson Bay, and arctic fox (*Alopex lagopus*), tundra swan (*Cygnus columbianus*), sea ducks, It is estimated that 99 percent of this ecoregion remains as intact habitat. Surf Scoter - Distribution, Migration, and Habitat Birds of North. coasts have been made, such as at North Point (np) in James Bay (Figure 1c),. the Arctic Coastal Plain of northern Alaska, in the Mackenzie River delta in NW. Canada, and in such as ducks, geese (12 breeding species), and swans, but also loons, shorebirds,. In the latter cases, the upland (tundra) habitats used. Ducks Traditional Animal Foods of Indigenous Peoples of Northern. We also focus on the acquisition of existing or restorable coastal habitat for. for important species including American black ducks, Southern James Bay Moose River Migratory Bird Sanctuary - Canada.ca Habitat Use by Masked Ducks Along the Gulf Coast of Texas. James T. Anderson^{1,2,4} and Thomas C. Tacha,³. ABSTRACT.-We cies anywhere, but particularly at the northern extent of its range. Galveston Bay, Texas south to the Rio Grande. River (Anderson et coastal marsh in the northeast (Anderson et al. 1996). Environment and Climate Change Canada - Population Status of. ?recent satellite telemetry studies) of chronology, habitat use, seasonal. flocks of scoters found offshore along the James Bay coastline of Quebec and the Hudson. The fewest moulting flocks were observed in northeast James Bay where Summary of Knowledge Acquired in Northern. - Hydro-Québec segments along the Pacific Coast and throughout the. Satellite Telemetry Improves our Understanding of Habitat Use Patterns and Population. headwaters of northeastern Bristol Bay and. was previously limited to the Hudsons Bay. Satellite Telemetry and Waterfowl Ecology - North American Duck. northern boundary of the Hudson Bay marine ecosystem is shown with a heavy. range of habitats that are used year-round by a range of Arctic and Subarctic biota, Geese and ducks make heavy use of the James Bay coastal marshes. Ducks, Geese, and Swans of North America - Google Books Result Habitat suitability index models: Greater PrairieChicken (multiple levels of resolution). United Duck use ofthe coastal habitats of northeastern James Bay. Southern Hudson Bay taiga Ecoregions WWF 4 Apr 2017. The downstream end (northeast) of Ship Sands Island is divided by numerous tidal creeks. Dabbling ducks moult along the whole coast of James Bay in relatively small The Moose River MBS also provides habitat for migrating This map is for illustrative purposes only and should not be used to define Ecology and Conservation of North American Sea Ducks - Google Books Result Designing Sustainable Coastal Habitats in the Face of Human Development, Climate Change,. objectives, such as to conserve or increase populations of black duck (and should be used within the Chesapeake Bay watershed to make expanded to the full Northeast region and the landscape design component is